GRASSES NEW TO EASTERN UNITED STATES

BY

CLYDE F. REED*

The following species of grasses seem to be new to the flora of Eastern United States. The specimens have been determined by Dr. Jason Swallen, Division of Grasses, United States National Herbarium. All the specimens are in the Reed Herbarium, unless noted otherwise.

Andropogon traceyi Nash South Carolina: Roadsides, Sharon, York Co. Dec. 26, 1955. Reed 37164.

Eragrostis oxylepis (Torr.) Torr. Virginia: Wastes at Fort Monroe, Hampton. Common in the sandy lawns. Elizabeth Co. Nov. 15, 1959. Reed 45919. Also in US.

<u>Festuca</u> <u>dertonensis</u> (All.) Aschers. & Graebn. Delaware: Edge of woods just east of Noxentown, New Castle Co. June 13, 1960. <u>Reed</u> 47196. Also in US.

Heleochloa schoeniodes (L.) Host. Delaware: Wharves, New Castle, New Castle Co. Oct. 21, 1863. <u>Albert Commons</u>. (Reed Herb.).

Many records for species of grasses new to Eastern United States were published by the author in 1964 (Phytologia, 10(5); 321-406) and 1965 (Phytologia, 11(5): 289).

* Reed Herbarium, 10105 Harford Road, Baltimore, Maryland, 21234.

